



Northwest Rheumatology Associates, P.C.
9555 SW Barnes Road - Suite 150 – Portland, Oregon 97225
Telephone: (503) 297-3384 - Fax: (503) 297-0863
www.nwrheumatology.org

Informed Consent for Treatment with Orenzia

Orenzia (Abatacept) is a medicine that is used to treat adults with moderate to severe Rheumatoid Arthritis (RA) who have not been helped by other medicines for RA. Orenzia keeps the immune system from attacking healthy tissues in the body. The immune system defends the body against infections caused by bacteria and viruses. A normal immune system leaves healthy body tissues alone. In people with RA, the immune system attacks normal body tissues causing damage and inflammation especially in the tissues of your joints. Orenzia interferes with an important step in this attack. By decreasing the immune system's attack on normal tissues, Orenzia can reduce pain and inflammation, and slow the damage to your bones and cartilage.

WARNINGS:

Orenzia decreases immune system function (immunosuppression), which can result in serious infections or worsening of an existing infection. Deaths from serious infections have been reported in patients receiving Orenzia.

Before starting treatment with Orenzia, you should have a tuberculosis skin test done to check for tuberculosis that may not be causing any symptoms (latent tuberculosis). If you have been diagnosed with tuberculosis, anti-tuberculosis medicines must be started before you start Orenzia therapy. This will reduce the likelihood of a serious tuberculosis infection. Tell your doctor if you have ever had tuberculosis, or have come into contact with someone with tuberculosis. Tell your doctor if you have ever had a positive PPD (tuberculosis skin test) – this may require treatment.

Tell your doctor immediately if you have any complaints that may indicate an infection. These include but are not limited to pink eye, ear pain or drainage, sinus pain or drainage, sore throat, hoarse voice, cough, difficulty breathing, fevers, chills, sweats, vomiting, abdominal pain, diarrhea, burning with urination, increased frequency of urination, vaginal discharge, and cuts or wounds that are red, swollen or draining pus.

Tell your doctor if you have been diagnosed as having an infection or if you are prone to recurrent infections. You should not receive Orenzia if you have an infection. You can receive Orenzia once your infection has been treated.

Tell your doctor or infusion nurse if you are on antibiotics around the time that your infusion or injection is due.

Tell your doctor if you have a planned surgery, as the timing of the surgery may need to be adjusted in accordance to your infusion/injection.

PRECAUTIONS:

Patients with a long duration of inflammation and chronic exposure to immunosuppressant treatments are more prone to develop cancer, in particular lymphoma. There have been rare cases of certain kinds of cancer in patients receiving orenicia. The role of orenicia in the development of cancer is not known.

Orenicia has not been studied in pregnant women or nursing mothers, so it is not known what the effects are on pregnant women or nursing babies. The safest approach at this time is to not use Orenicia if you are pregnant, breast feeding, become pregnant, or are thinking about becoming pregnant.

You should not receive live vaccines (for eg. polio, smallpox) while on Orenicia or within 3 months of having received Orenicia. The injected flu vaccine and pneumonia vaccine can be safely administered while on Orenicia. The nasal spray version of the flu vaccine is a live vaccine and should not be administered while on Orenicia. Orenicia may blunt the effectiveness of some vaccines. No data is available on the transmission of infection from persons receiving live vaccines to persons receiving Orenicia. Please let your doctor know if someone in your household has received or will receive a live virus vaccine.

ADVERSE REACTIONS:

Infusion related reactions have occurred in about 9% of patients treated with Orenicia in clinical trials.

These reactions include but are not limited to headache, high blood pressure, dizziness, low blood pressure, shortness of breath, nausea, flushing, hives, cough, hypersensitivity, itching, rash, wheezing, and rarely anaphylaxis. Most of these reactions were mild to moderate, and responded to appropriate treatment, or to slowing down or stopping the infusion.

Injection site reactions can occur resulting in redness or itching at the site of the injection.

Patients with COPD (chronic obstructive pulmonary disease, emphysema) developed adverse events more frequently. These adverse events included COPD exacerbations, cough, wheezing, and shortness of breath. Tell your doctor if you have COPD or if your breathing worsens in any way.

MEDICATION ADMINISTRATION:

Orenicia is administered as a single intravenous infusion lasting about 30 minutes. You will receive your first dose of Orenicia followed by additional doses at 2 and 4 weeks after the first dose. You will then receive a dose every 4 weeks. You will see your physician in follow-up to monitor your progress and response to Orenicia.

Orenicia is also administered as a subcutaneous injection every week.

Please call us if you experience any side effects, and we will be available to you 24 hours a day if you experience any problems. We may give you additional medication to take before the infusion to reduce the risk of side effects. If you receive these medications, you may need someone to drive you home after the infusion.

EMERGENCY CONTACTS:

In case of emergency, you can contact our office at (503) 297-3384 from 8:00 am to 5:00 pm Monday – Friday, or after hours dial the above number to page your physician or call 911 or go to the emergency room.

I certify that I have read and understand this consent form and agree to receive the Orencia intravenous treatment. I have had an opportunity to discuss this treatment with my physician and ask questions regarding the treatment. I will be given a signed copy of this form for my records.

Patient signature

Date

FINANCIAL RESPONSIBILITY:

Our office will contact your insurance company and the necessary arrangements will be made for approval of the medication and administration. You may want to follow up with your insurance company also to be sure everything is covered. **Our office does NOT provide a guarantee of coverage for any of these services. If coverage is not provided, YOU WILL BE RESPONSIBLE for any charges incurred for treatment and/or follow-up care.**

Patient signature

Date